Initial REMS approval: 02/2014

Most recent modification: 06/2014

BLA 125390 MYALEPT™ (metreleptin)

Drug Class: Recombinant human leptin analog

Manufactured for Amylin Pharmaceuticals LLC¹
1800 Concord Pike
P.O Box 8355
Wilmington, DE, 19803-8355
1-800-236-9933

MYALEPT is marketed and distributed by: AstraZeneca Pharmaceuticals LP 1800 Concord Pike
P.O Box 8355
Wilmington, DE, 19803-8355
1-800-236-9933

RISK EVALUATION AND MITIGATION STRATEGY (REMS)

I. GOALS

The goal of the MYALEPT REMS is to mitigate (1) the risks of serious adverse sequelae (such as severe infections, excessive weight gain, glucose intolerance, diabetes mellitus) due to the development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT, and (2) the risk of lymphoma by:

- Educating prescribers about the development of neutralizing anti-metreleptin antibodies, the serious adverse sequelae that may result from these antibodies, and the risk for lymphoma associated with MYALEPT
- Limiting the population exposed to MYALEPT by requiring prescriber certification, pharmacy certification, and prescriber attestation that each patient has a diagnosis consistent with the approved indication

Reference ID: 3534419

¹ Amylin Pharmaceuticals, LLC (Amylin Pharmaceuticals) is the licensee under BL 125390. Although Amylin Pharmaceuticals and AstraZeneca Pharmaceuticals LP have a contractual agreement under which AstraZeneca Pharmaceuticals LP is to carry out the responsibilities under the REMS for MYALEPT on behalf of Amylin Pharmaceuticals, Amylin Pharmaceuticals retains primary responsibility for all actions described in the REMS.

II. REMS ELEMENTS

A. Elements to Assure Safe Use (ETASU)

- 1. Healthcare Providers (HCPs) who prescribe MYALEPT are specially certified.
 - a. Amylin Pharmaceuticals will ensure that HCPs who prescribe MYALEPT are specially certified.
 - b. To become specially certified to prescribe MYALEPT, prescribers must enroll in the MYALEPT REMS Program. Prescribers must complete the following requirements:
 - i. review the Prescribing Information
 - ii. complete the Prescriber Training Module
 - iii. complete and sign the *Prescriber Enrollment Form* and submit it to the MYALEPT REMS Program.
 - c. Amylin Pharmaceuticals will:
 - i. Ensure that the *Prescriber Training Module* and the *Prescriber Enrollment Form* are available on the MYALEPT REMS Program website (www.MYALEPTREMS.com) or can be obtained by contacting the MYALEPT REMS Program by phone at 1-855-6MYALEPT.
 - ii. Ensure that prescribers complete the *Prescriber Training Module* and the *Prescriber Enrollment Form* before activating prescribers' certification in the MYALEPT REMS Program.
 - iii. Ensure that prescribers are notified when they have been successfully certified by the MYALEPT REMS Program.
 - iv. Inform certified prescribers following substantial changes to the MYALEPT REMS or MYALEPT REMS Program; substantial changes include significant changes to the operation of the MYALEPT REMS Program or changes to the Prescribing Information that affect the risk-benefit profile of MYALEPT.
 - v. Communicate information about the risks and MYALEPT REMS Program requirements to prescribers through the MYALEPT REMS Program website and MYALEPT REMS introductory information sheet titled *MYALEPT REMS Program: An Introduction*. Amylin Pharmaceuticals will provide the *REMS Program: An Introduction, Prescriber Training Module, Prescriber Enrollment Form*, and the Prescribing Information to prescribers who (1) attempt to prescribe MYALEPT and are not yet certified, or (2) inquire about how to become certified.

The following materials are part of the REMS and are appended:

- Prescriber Training Module
- Prescriber Enrollment Form
- MYALEPT REMS Program: An Introduction

• MYALEPT REMS Program website

2. Pharmacies that dispense MYALEPT are specially certified.

a. Amylin Pharmaceuticals will ensure that MYALEPT is dispensed only by specially certified pharmacies.

To become certified to dispense MYALEPT, each pharmacy representative must agree to the following:

- i. to educate all pharmacy staff involved in the dispensing of MYALEPT on the MYALEPT REMS Program requirements
- ii. to put processes and procedures in place to verify, prior to dispensing MYALEPT, that:
 - 1) the prescriber is certified in the MYALEPT REMS Program
 - 2) the *MYALEPT REMS Prescription Authorization Form* is received for each new prescription
- iii. to be audited to ensure that all processes and procedures are in place and are being followed for the MYALEPT REMS Program
- iv. to provide prescription data to the MYALEPT REMS Program, and
- v. to refrain from reselling or transferring MYALEPT to other pharmacies or distributors.

The following material is part of the REMS and appended:

• MYALEPT REMS Prescription Authorization Form

3. MYALEPT is dispensed only to patients with evidence or other documentation of safe-use conditions.

- a. MYALEPT is dispensed only to patients whose prescribers are specially certified in the MYALEPT REMS Program and attest on the MYALEPT REMS Prescription Authorization Form that:
 - i. They understand that MYALEPT is indicated as an adjunct to diet as replacement therapy to treat the complications of leptin deficiency in patients with congenital or acquired generalized lipodystrophy.
 - ii. They affirm that their patient has a clinical diagnosis consistent with the approved indication, and that the patient (or caregiver) has been properly informed of the benefits and risks of MYALEPT therapy.
 - iii. They understand that MYALEPT is not indicated for the treatment of complications of partial lipodystrophy.
 - iv. They understand that MYALEPT is not indicated for the treatment of liver disease, including non-alcoholic steatohepatitis (NASH).
 - v. They understand that MYALEPT is not indicated for use in patients with HIV-related lipodystrophy.
 - vi. They understand that MYALEPT is not indicated for use in patients with metabolic disease, including diabetes mellitus and hypertriglyceridemia, without concurrent evidence of congenital or acquired generalized lipodystrophy.
 - vii. They understand that MYALEPT is contraindicated in patients with general obesity not associated with congenital leptin deficiency.
 - viii. They understand that MYALEPT is associated with serious adverse events due to the development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT.
 - ix. They agree to test for neutralizing antibodies to MYALEPT if a patient experiences severe infections or if they suspect MYALEPT treatment is no longer working (e.g., loss of glycemic control, increased triglycerides).
 - x. They understand that MYALEPT is associated with a risk of lymphoma.

xi. They understand it is important to carefully consider treatment with MYALEPT in patients with significant hematologic abnormalities and/or acquired generalized lipodystrophy

B. Implementation System

- 1. Amylin Pharmaceuticals will ensure that MYALEPT is distributed to and dispensed only by certified pharmacies.
- 2. Amylin Pharmaceuticals will maintain, monitor, and evaluate the implementation of the MYALEPT REMS Program.
 - a. Amylin Pharmaceuticals will maintain a secure, validated database of certified prescribers and pharmacies in the MYALEPT REMS Program. Amylin Pharmaceuticals will send confirmation of certification to each certified pharmacy.
 - b. Amylin Pharmaceuticals will maintain a MYALEPT REMS Program Call Center to support patients, prescribers, and pharmacies interfacing with the MYALEPT REMS.
 - c. Amylin Pharmaceuticals will ensure that all materials listed in or appended to the MYALEPT REMS Program will be available through the MYALEPT REMS Program website (www.MYALEPTREMS.com) or by calling the MYALEPT REMS Program Call Center at 1-855-6MYALEPT.
 - d. If there are substantive changes to the MYALEPT REMS or MYALEPT REMS Program, Amylin Pharmaceuticals will update all affected materials, and notify enrolled prescribers and certified pharmacies, as applicable. Substantive changes are defined as significant changes to the operation of the MYALEPT REMS Program or changes to the Prescribing Information that affect the risk-benefit profile of MYALEPT.
 - e. Amylin Pharmaceuticals will monitor and audit the certified pharmacies within 180 days after the pharmacy is certified to ensure that all processes and procedures are in place and functioning to support the requirements of the MYALEPT REMS Program. Thereafter, Amylin Pharmaceuticals will include the certified pharmacies in the company's annual audit plan. Corrective action will be instituted by Amylin Pharmaceuticals if noncompliance is found.
 - f. Based on monitoring and evaluation of the MYALEPT REMS elements to assure safe use, Amylin Pharmaceuticals will take reasonable steps to improve implementation of these elements and to maintain compliance with the MYALEPT REMS Program requirements, as applicable.

C. Timetable for Submission of Assessments

Amylin Pharmaceuticals will submit REMS Assessments to the FDA 6 months, 12 months, and annually thereafter from the date of the initial REMS approval. To facilitate inclusion of as much information as possible while allowing reasonable time to prepare the submission, the reporting

interval covered by each assessment should conclude no earlier than 60 days before the submission date for that assessment. Amylin Pharmaceuticals will submit each assessment so that it will be received by the FDA on or before the due date.



MYALEPT[™] REMS Program Prescriber Enrollment Form

11.3 mg per vial

MYALEPT will be available only through the MYALEPT REMS Program. To prescribe MYALEPT, a prescriber must: (1) review the Prescribing Information, review and complete the Prescriber Training Module, (2) complete this one-time MYALEPT REMS Program Prescriber Enrollment Form, and (3) complete and submit a MYALEPT REMS Prescription Authorization Form for each new prescription.

Complete this enrollment form and	d fax it to the MYALE		gram	at 1-877	-328-9682.	
Prescriber Information (Please P	rint *indicates a red	quired field.)				
Full Name (first, middle, last)*						
Credentials*						
□MD □DO □NP □PA □O	ther (specify)					
Physician Specialty*	☐ Family Mo Medicine ☐ Pediatrics		ardiolog ther	gy		
Who do you treat?						
☐ Adults ☐ Pediatrics ☐ Both						
Practice / Facility Name						
Address 1*						
Address 2 (optional)		City*			State*	ZIP code*
Phone number*	Alternate phone number*			Fax numbe	<u> </u> 	
Email*		NPI #*				
Practice Setting Solo private practice			emic/ I	Hospital a	affiliated pra	ctice
Office Contact						
Full Name (first, middle, last)*						
Phone number (if different from above) Fax no	umber (if different from abov	re) Email* (if office	ce conta	ct is provide	ed)	
Prescriber Attestation. By completing the I understand that MYALEPT is indicated as a	n adjunct to diet as replace	ement therapy to trea	at the co	omplications	s of leptin-defi	ciency in patients
with congenital or acquired generalized lipod I understand that MYALEPT is available only	ystrophy. through the MYALEPT RE	MS Program and the	at I mus	st comply w	ith the prograr	n requirements in
order to prescribe MYALEPT. I have completed the MYALEPT REMS Pres	oribor Training Madula	-				
I understand that MYALEPT is associated with the state of the sta		due to the developme	ent of a	nti-metreler	otin antibodies	that neutralize
endogenous leptin and/or MYALEPT.		·				
 I agree to test for neutralizing antibodies in p glycemic control, or increases in triglycerides 	atients wno experience sev :).	ere intections or it is	suspect	WYALEPI	is no longer v	vorking (e.g., loss of
 I understand that MYALEPT is associated wi 	th a risk of lymphoma.				,	
 I will carefully consider the risks of treatment lipodystrophy. 	with MYALEP1 in patients	with significant hema	atologic	abnormalit	ies and/or acc	luired generalized
 I understand that MYALEPT is not indicated 	for the treatment of complic	ations of partial lipod	dystropl	hy.		
I understand that MYALEPT is <u>not</u> indicated				lic steatohe	patitis (NASH)).
 I understand that MYALEPT is <u>not</u> indicated I understand that MYALEPT is not indicated 	for use in patients with HIV- for use in patients with met	-related lipodystroph abolic disease, inclu	1y. Idina dis	hetes melli	tus and hyner	trialyceridemia
without concurrent evidence of congenital or	acquired generalized lipody	ystrophy.	ŭ		•	•
 I understand that MYALEPT is contraindicate I agree that personnel from the MYALEPT R 	EMS Program may contact	me to gather further	ed with o	congenital le ation or res	eptin deficienc olve discrepar	y. ncies or to provide
 other information related to MYALEPT or the I will complete and submit a MYALEPT REM I agree that AstraZeneca, its agents and con the effectiveness of the program requirement 	S Program Prescription Aut tractors such as the pharma	thorization Form for acy providers, may d	each ne contact i	ew prescript me via phoi	tion. ne, mail, or en	nail to survey me on
Signature*	13 IOI IIIE IVITALEET EREIVIS	i iogiaiii.		Date*		

If you have any questions, please contact the MYALEPT REMS Program.

Phone number: 1-855-669-2537 | Fax number: 1-877-328-9682 | www.MYALEPTREMS.COM

AstraZeneca



MYALEPT[™] REMS Program **Prescription Authorization Form**

Instructions: Complete both **pages** of this form for each new prescription. All fields are required. Please Print. Please FAX completed form to MYALEPT REMS Program at 1-877-328-9682.

The prescription for MYALEPT is only valid if received by fax.

For New York prescribers: In addition to this completed form, provide New York specific prescription blanks.

Patient Information				
Full Name (first, middle, last)		Gender □ Male □ Fema	le Date of Birth	
Address		City	State	ZIP code
Preferred phone number	Alternate phone nun	nber	Preferred time to c □ Day □ Eveni	ontact: (check one) ng
Email		Alternate contact/phone #		
Parent/Guardian (if applicable)				
Insurance Information	Please copy and atta	ach the front and bac	k of the insura	ince card.
Insurance company phone number				
Insured Name		Relationship to patient		
Insured Employer				
Prescription card ☐ Yes ☐ N	o If yes, carrier			
Policy Number				
Is the patient eligible for Medicare?	□ Yes □ No			
Policy number		Group Number		
Shipping Information				
Full Name (first, middle, last)				
Address (if different from above)		City	State	ZIP code
Send initial shipment to prescribing	doctor's office ☐ Yes ☐ No			
Prescriber Information				
Full Name (first, middle, last)				
Practice / Facility Name		Office Contact Person		
Address 1				
Address 2		City	State	ZIP code
Office Phone number	Office Fax number	License #	NPI#	

Continue to page 2 to complete prescription.

The MYALEPT REMS Program may be found at www.MYALEPTREMS.com or by calling 1-855-669-2537.





Physician Signature: _

11.3 mg per vial

MYALEPT[™] REMS Program Prescription Authorization Form

Attestation of REMS Requirements. By completing this form, I attest that:

- I understand that MYALEPT is indicated as an adjunct to diet as replacement therapy to treat the complications of leptin-deficiency in patients with congenital or acquired generalized lipodystrophy.
- I affirm that my patient has a clinical diagnosis consistent with generalized lipodystrophy, and that my patient (or their caregiver) has been properly informed of the benefits and risks of MYALEPT therapy.
- I understand that MYALEPT is <u>not</u> indicated for the treatment of complications of partial lipodystrophy.
- I understand that MYALEPT is not indicated for the treatment of liver disease, including non-alcoholic steatohepatitis (NASH).
- I understand that MYALEPT is not indicated for use in patients with HIV-related lipodystrophy.
- I understand that MYALEPT is <u>not</u> indicated for use in patients with metabolic disease including diabetes mellitus and hypertriglyceridemia without concurrent evidence of congenital or acquired generalized lipodystrophy.
- I understand that MYALEPT is contraindicated in patients with general obesity not associated with congenital leptin deficiency.
- I understand that MYALEPT is associated with serious adverse events due to the development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT.
- I agree to test for neutralizing antibodies in patients who experience severe infections or if I suspect MYALEPT is no longer working (e.g., loss of glycemic control, or increases in triglycerides).
- I understand that MYALEPT is associated with a risk of lymphoma.
- I understand I must carefully consider the risks of treatment with MYALEPT in patients with significant hematological abnormalities and/or acquired generalized lipodystrophy.

Starting Dose: [] 0.06 mg/kg [] 2.5 mg [] 5 mg [Patient Weight Days Supply		Refills #		
Directions (e.g., by subcutaneous injection once daily)		IVEIIIIS #		
List or Attach a List of Concomitant Medications			Allergies	
			J	
Full Name (first, middle, last)				
Full Name (first, middle, last)				
Address				
City	State		ZIP code	Office Phone
License #			NPI#	·
hysician Signature:	Date	Phy	sician Signaturo:	Date
Product Selection Permitted	Date	FIIY	oronam orginature.	Dispense as Written
YALEPT is manufactured for Amylin Pharmaceuticals and dis-	stributed by A	ry supplies fo	r MYALEPT reco	onstitution and administration. This can be given to your patient to fill at another phan
YALEPT is a trademark of he AstraZeneca group of compar YALEPT is manufactured for Amylin Pharmaceuticals and discrete following is a prescription form for the require ack to 1-877-328-9682 with the rest of this page Patient Information	stributed by A	ry supplies fo	r MYALEPT reco	onstitution and administration. This can be
YALEPT is manufactured for Amylin Pharmaceuticals and discrete following is a prescription form for the require ack to 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last)	stributed by A	ry supplies fo	r MYALEPT reco	onstitution and administration. This can be
YALEPT is manufactured for Amylin Pharmaceuticals and discrete following is a prescription form for the require ack to 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last) MYALEPT Supplies Prescription	ed ancillar e and the p	ry supplies fo previous page	r MYALEPT reco	onstitution and administration. This can be given to your patient to fill at another phan
YALEPT is manufactured for Amylin Pharmaceuticals and discher following is a prescription form for the require ack to 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last) MYALEPT Supplies Prescription Required supplies (please note - the maximum numb	ed ancillar e and the p	ry supplies fo previous page	r MYALEPT reco	onstitution and administration. This can be given to your patient to fill at another phane
YALEPT is manufactured for Amylin Pharmaceuticals and discher following is a prescription form for the require ack to 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last) MYALEPT Supplies Prescription	ed ancillar e and the p	ry supplies fo previous page	oelow. Pharmacy v	onstitution and administration. This can be given to your patient to fill at another phane
PALEPT is manufactured for Amylin Pharmaceuticals and discrete following is a prescription form for the requirement to 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last) MYALEPT Supplies Prescription Required supplies (please note - the maximum numb for Reconstitution 5 of 30 mL vials of BWFI Refills # 31 of 5 mL vials of SWFI (use for neonates and infants Refills # 62 of 3mL syringe (22G x 1 in. needle) Refills # 1 decomposition 1 decomp	ed ancillar e and the p	ry supplies fo previous page	oelow. Pharmacy v	onstitution and administration. This can be given to your patient to fill at another phane will adjust to individual patient needs.) inistration tuberculin syringe (26 G x 3/8 in. needle)
ALEPT is manufactured for Amylin Pharmaceuticals and discrete following is a prescription form for the require ack to 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last) MYALEPT Supplies Prescription Required supplies (please note - the maximum numb for Reconstitution 5 of 30 mL vials of BWFI Refills # 31 of 5 mL vials of SWFI (use for neonates and infants Refills # 62 of 3mL syringe (22G x 1 in. needle) Refills # Prescriber Information	ed ancillar e and the p	ry supplies fo previous page	oelow. Pharmacy v	onstitution and administration. This can be given to your patient to fill at another phane will adjust to individual patient needs.) inistration tuberculin syringe (26 G x 3/8 in. needle)
Packet of 1-877-328-9682 with the rest of this page Patient Information Required supplies (please note - the maximum numbor of 30 mL vials of BWFI Refills #	ed ancillar e and the p	ry supplies fo previous page	oelow. Pharmacy v	onstitution and administration. This can be given to your patient to fill at another phane will adjust to individual patient needs.) inistration tuberculin syringe (26 G x 3/8 in. needle)
Patient Information Full Name (first, middle, last) Refills # 31 of 5 mL vials of SWFI (use for neonates and infants Refills # 62 of 3mL syringe (22G x 1 in. needle) Refills # Prescriber Information Full Name (first, middle, last)	ed ancillar e and the p	ry supplies fo previous page	oelow. Pharmacy v	onstitution and administration. This can be given to your patient to fill at another phane will adjust to individual patient needs.) inistration tuberculin syringe (26 G x 3/8 in. needle)
Packet of 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last) WYALEPT Supplies Prescription Required supplies (please note - the maximum numb For Reconstitution 5 of 30 mL vials of BWFI Refills # 31 of 5 mL vials of SWFI (use for neonates and infants Refills # 62 of 3mL syringe (22G x 1 in. needle) Refills # Prescriber Information Full Name (first, middle, last)	ed ancillar e and the per supp	ry supplies fo previous page	pelow. Pharmacy v For Admir G2 of 1mL Refills #_	onstitution and administration. This can be given to your patient to fill at another phane will adjust to individual patient needs.) nistration tuberculin syringe (26 G x 3/8 in. needle)
YALEPT is manufactured for Amylin Pharmaceuticals and discrete following is a prescription form for the require ack to 1-877-328-9682 with the rest of this page Patient Information Full Name (first, middle, last) MYALEPT Supplies Prescription Required supplies (please note - the maximum number for Reconstitution 5 of 30 mL vials of BWFI (use for neonates and infants Refills #	ed ancillar e and the per per supp	ry supplies fo previous page oly is specified	r MYALEPT recorder, or torn off and poelow. Pharmacy via the second of t	onstitution and administration. This can be given to your patient to fill at another phane will adjust to individual patient needs.) nistration tuberculin syringe (26 G x 3/8 in. needle)




11.3 mg per vial

Risk Evaluation and Mitigation Strategy (REMS) Program
Prescriber Training Module

Contents

- Introduction
- MYALEPT[™] (metreleptin) for injection Product Information
 - Boxed Warning: risk of development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT
 - Boxed Warning: risk of lymphoma
 - Appropriate patient selection
- MYALEPT REMS Program Information
- Knowledge Assessment



Introduction

MYALEPT™ (metreleptin) for injection is available only through a restricted program called the MYALEPT REMS Program.

 Prescribers must complete this training module and enroll in the MYALEPT REMS Program prior to prescribing MYALEPT.

The purpose of this training module is to educate prescribers about

- the development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT and the serious adverse events that may result from these antibodies,
- the risk of lymphoma, and
- appropriate patient selection

Because of these risks, appropriate patient selection consistent with the approved indication for MYALEPT is very important.



Indication

MYALEPT™ (metreleptin) for injection is indicated as an adjunct to diet as replacement therapy to treat the complications of leptin deficiency in patients with congenital or acquired generalized lipodystrophy

Serious Risks Associated with MYALEPT

Serious Risks Associated with MYALEPT

- Development of anti-metreleptin antibodies that neutralize endogenous leptin and MYALEPT
- 2. Lymphoma

Boxed Warning: Anti-metreleptin Antibodies with Neutralizing Activity - Serious Adverse Events

- Anti-metreleptin antibodies with neutralizing activity associated with adverse events consistent with loss of endogenous leptin activity and/or loss of efficacy have been identified among patients with generalized lipodystrophy
 - 2 of 33 patients who underwent antibody testing
 - tested positive for anti-metreleptin antibodies with neutralizing activity
 and
 - reported adverse events consistent with neutralizing activity, including severe infections, loss of glycemic control, and increases in triglycerides

Boxed Warning: Anti-metreleptin Antibodies with Neutralizing Activity - Serious Adverse Events (continued)



In other populations

- 3 of 563 patients who underwent antibody testing
 - tested positive for anti-metreleptin antibodies with neutralizing activity
 and
 - reported adverse events consistent with neutralizing activity, including excessive weight gain and development of glucose intolerance or diabetes mellitus

The clinical implications associated with development of anti-metreleptin antibodies with neutralizing activity are not well-characterized at this time due to the small number of reports.

Antibody Testing in MYALEPT Trials - Data

- Anti-metreleptin antibodies were detected in 84% (36/43) of patients with generalized lipodystrophy studied in the MYALEPT trials
- Total anti-metreleptin antibody titers ranged between 1:5 and 1:1,953,125
- Anti-metreleptin antibodies with neutralizing activity associated with adverse events consistent with loss of endogenous leptin activity and/or loss of MYALEPT efficacy were observed in 6% (2/33) of the patients tested

Antibody Testing in MYALEPT Trials - Limitations

- The immunogenicity assays utilized in clinical trials lacked sensitivity, resulting in potential underestimation of the number of samples positive for anti-metreleptin antibodies with neutralizing activity
 - The observed incidence of an antibody assay (including neutralizing antibody assays) positivity may be influenced by several factors including assay sensitivity and specificity, assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease.

Antibody Testing in MYALEPT Trials – Limitations (continued)

- Antibody testing was not performed uniformly in the MYALEPT trials. The incompleteness of the current immunogenicity database precludes understanding of the magnitude and persistence of the observed anti-metreleptin antibody responses.
- Comparison of the incidence of antibodies to metreleptin with the incidence of antibodies to other products may be misleading.

Neutralizing Activity - What is My Role?

- Test for neutralizing activity in patients who experience severe infections or if you suspect that MYALEPT is no longer working.
- Contact AstraZeneca at 1-866-216-1526 for instructions on how to submit samples for neutralizing antibody testing. The assay is not commercially available.

AstraZeneca will ask you to:

- 1. Obtain written consent from your patient to release the sample and send a copy of the consent to AstraZeneca.
- 2. Complete a questionnaire to explain why you are requesting neutralizing activity testing and send it to AstraZeneca.
- 3. Send the sample to the designated laboratory for testing.
 - The results are generally available within 60 days
- 4. AstraZeneca will contact you to provide and discuss the results.

Boxed Warning: Lymphoma

- Three cases of T-cell lymphoma have been reported in the MYALEPT lipodystrophy program
 - All 3 patients had acquired generalized lipodystrophy (out of a total of 20 patients with acquired generalized lipodystrophy).
 - Two of these patients were diagnosed with peripheral T-cell lymphoma while receiving MYALEPT.
 - Both had immunodeficiency and significant hematologic abnormalities, including severe bone marrow abnormalities, before the start of MYALEPT treatment.
 - A separate case of anaplastic large cell lymphoma was reported in a patient receiving MYALEPT who did not have hematological abnormalities before treatment.

Lymphoma – What is My Role?

- Take a careful medical history for past or current hematologic abnormalities
- Carefully consider the benefits and risks of treatment with MYALEPT in patients with
 - > significant hematologic abnormalities, and/or
 - acquired generalized lipodystrophy

Adverse Reaction Reporting

To report SERIOUS ADVERSE REACTIONS, please call/contact:

- 1-855-669-2537 and/or
- FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

Appropriate Patient Selection

Appropriate Patient Selection

Contraindication – General Obesity

MYALEPT is contraindicated in patients with general obesity not associated with congenital leptin deficiency.

- MYALEPT has not been shown to be effective in treating general obesity, and the development of anti-metreleptin antibodies with neutralizing activity has been reported in obese patients treated with MYALEPT
- Adverse events consistent with loss of endogenous leptin activity have been identified in three patients <u>without</u> lipodystrophy who received metreleptin (excessive weight gain, development of glucose intolerance or diabetes mellitus).

The clinical implications associated with development of anti-metreleptin antibodies with neutralizing activity are not well-characterized at this time due to the small number of reports.

Appropriate Patient Selection

Important Limitations of Use

- The safety and effectiveness of MYALEPT for the following conditions have <u>not</u> been established
 - The treatment of complications of partial lipodystrophy
 - The treatment of liver disease including non-alcoholic steatohepatitis (NASH)
- MYALEPT is not indicated for use in patients with
 - HIV-related lipodystrophy
 - Metabolic disease including diabetes mellitus and hypertriglyceridemia, without concurrent evidence of congenital or acquired generalized lipodystrophy.

MYALEPT™ (metreleptin) for injection REMS PROGRAM INFORMATION

MYALEPT™ (metreleptin) for injection REMS Program



Key program elements:

- Certification of Prescribers of MYALEPT
 - Certification consists of training, and enrolling in the MYALEPT REMS Program.
- Completion of a Prescription Authorization Form for each new prescription
- Restricted distribution of MYALEPT through certified pharmacies.

MYALEPT™ (metreleptin) for injection REMS Program

Before prescribing MYALEPT, prescribers must complete the following steps:

- Review the Prescribing Information and this Prescriber Training Module.
- Complete, sign, and submit the one-time MYALEPT REMS Program Prescriber Enrollment Form.
- Complete, sign, and submit the MYALEPT REMS Program Prescription Authorization Form for each new prescription.

Note: All Materials can be downloaded from the MYALEPT REMS website at: www.MYALEPTREMS.com

Or request these materials by calling 1-855-669-2537

1. Review Prescriber Education Materials

Review the following Prescriber Education Materials:

- a) MYALEPT™ (metreleptin) for injection Prescribing Information
- b) This Prescriber Training Module

2. Enroll in MYALEPT™ (metreleptin) for injection REMS Program

To enroll in the MYALEPT REMS Program:

- Download the MYALEPT REMS Program Prescriber Enrollment Form at www.MYALEPTREMS.com or request a copy by calling 1-855-669-2537
- Complete the enrollment form
- Sign & submit the enrollment form
 - Fax to 1-877-328-9682

3. Submit Prescription Authorization Form

When prescribing MYALEPT™ (metreleptin) for injection, a prescriber must complete a Prescription Authorization Form for each new prescription.

As part of completing the Prescription Authorization Form, you attest that:

- I understand that MYALEPT is indicated as an adjunct to diet as replacement therapy to treat the complications of leptin-deficiency in patients with congenital or acquired generalized lipodystrophy.
- I affirm that my patient has a clinical diagnosis consistent with generalized lipodystrophy, and that my patient (or their caregiver) has been properly informed of the benefits and risks of MYALEPT therapy.

Submit Prescription Authorization Form (continued)



- I understand that MYALEPT is <u>not</u> indicated for the treatment of complications of partial lipodystrophy.
- I understand that MYALEPT is <u>not</u> indicated for the treatment of liver disease, including non-alcoholic steatohepatitis (NASH).
- I understand that MYALEPT is not indicated for use in patients with HIV-related lipodystrophy.
- I understand that MYALEPT is <u>not</u> indicated for use in patients with metabolic disease including diabetes mellitus and hypertriglyceridemia without concurrent evidence of congenital or acquired generalized lipodystrophy.

Submit Prescription Authorization Form (continued)

- I understand that MYALEPT is contraindicated in patients with general obesity not associated with congenital leptin deficiency.
- I understand that MYALEPT is associated with serious adverse events due to the development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT.
- I agree to test for neutralizing antibodies in patients who experience severe infections or if I suspect MYALEPT is no longer working (e.g., loss of glycemic control, or increases in triglycerides).
- I understand that MYALEPT is associated with a risk of lymphoma.
- I understand I must carefully consider the risks of treatment with MYALEPT in patients with significant hematologic abnormalities and/or acquired generalized lipodystrophy.


3. Submit Prescription Authorization Form



- Download the Prescription Authorization Form at www.MYALEPTREMS.com or request a copy by calling 1-855-669-2537
- 2. Complete the Prescription Authorization Form
- 3. Sign & submit the Prescription Authorization Form
 - Fax to 1-877-328-9682



- The following questions about MYALEPT™ (metreleptin) for injection are provided to reinforce learning.
- If you have difficulty answering these questions, review the previous slides and refer to the Prescribing Information.




with congenital or acquired generalized lipodystrophy.

- 1. Which of the following statements is true?
 - MYALEPT (metreleptin) is indicated for use in patients with HIV-related lipodystrophy.
 - MYALEPT is indicated for use in patients with metabolic disease, including diabetes mellitus and hypertriglyceridemia, without concurrent evidence of inherited or acquired generalized lipodystrophy.
 - ☐ MYALEPT is a recombinant analog of murine leptin.
 - ✓ MYALEPT is indicated as an adjunct to diet as replacement therapy to treat the complications of leptin deficiency in patients with congenital or acquired generalized lipodystrophy.

Answer: MYALEPT is indicated as an adjunct to diet as replacement therapy to treat the complications of leptin deficiency in patients with congenital or acquired generalized lipodystrophy

(continued on next slide)



- The safety and effectiveness of MYALEPT for the following conditions have not been established
 - The treatment of complications of partial lipodystrophy
 - The treatment of liver disease including non-alcoholic steatohepatitis (NASH)
- MYALEPT is not indicated for use in patients with
 - HIV-related lipodystrophy
 - Metabolic disease including diabetes mellitus and hypertriglyceridemia, without concurrent evidence of congenital or acquired generalized lipodystrophy.
- MYALEPT is contraindicated in patients with general obesity not associated with congenital leptin deficiency.

The risks and benefits of MYALEPT™ (metreleptin) for injection treatment should be carefully considered in patients with significant hematologic abnormalities (for example, leukopenia, neutropenia, bone marrow abnormalities, lymphoma and/or lymphadenopathy) and/or acquired generalized lipodystrophy.

☐ True

☐ False

- The risks and benefits of MYALEPT™ (metreleptin) for injection treatment should be carefully considered in patients with significant hematologic abnormalities (for example, leukopenia, neutropenia, bone marrow abnormalities, lymphoma and/or lymphadenopathy) and/or acquired generalized lipodystrophy.
 - ✓ True
 - ☐ False
- Answer: Peripheral T-cell lymphoma was diagnosed in two patients with acquired generalized lipodystrophy while receiving MYALEPT.
 - Both had immunodeficiency and significant hematologic abnormalities including severe bone marrow abnormalities before the start of MYALEPT treatment.
 - A separate case of anaplastic large cell lymphoma was reported in a patient with acquired generalized lipodystrophy who did not have hematologic abnormalities before MYALEPT treatment.

- 3. Developing neutralizing activity to metreleptin could:
 - ☐ affect endogenous leptin
 - ☐ result in loss of efficacy
 - □ both of the above
 - none of the above

- 3. Developing neutralizing activity to metreleptin could:
 - ☐ affect endogenous leptin
 - result in loss of efficacy
 - ✓ both of the above
 - none of the above

Answer: Developing neutralizing activity to metreleptin could affect endogenous leptin and could result in loss of efficacy.

4. If a patient is experiencing severe infections and/or I suspect Myalept is no longer working, I will contact AstraZeneca for instructions on how to send a blood sample to test for anti-metreleptin antibodies with neutralizing activity.

☐ True

☐ False

- 4. If a patient is experiencing severe infections and/or I suspect Myalept is no longer working, I will contact AstraZeneca for instructions on how to send a blood sample to test for anti-metreleptin antibodies with neutralizing activity.
 - ✓ True
 - False

Answer: If you suspect your patient is experiencing complications from the development of anti-metreleptin neutralizing antibodies, you can submit a request and obtain results from AstraZeneca at no cost.

- 1. Call 1-866-216-1526
 - AstraZeneca will provide you information on the requirements for sample collection and shipment.
- AstraZeneca will instruct you to:
 - a. Obtain written consent from your patient to release the sample and send a copy of the consent to AstraZeneca.
 - b. Complete a questionnaire to explain why you are requesting neutralizing activity testing.
 - c. Send the sample and paperwork to the designated laboratory for testing.
 - The results are generally available within 60 days
- AstraZeneca will contact you to provide and discuss the results.

- 5. How often should the Prescription Authorization Form be completed?
 - ☐ Each new prescription
 - ☐ Only on the first prescription
 - ☐ Every refill
 - Once a year

- 5. How often should the Prescription Authorization Form be completed?
 - ✓ Each new prescription
 - Only on the first prescription
 - Every refill
 - Once a year

Answer: For each new prescription, the prescriber must submit a Prescription Authorization Form.



□ True

☐ False

- MYALEPT™ (metreleptin) for injection is available only through certified pharmacies.
 - ✓ True
 - ☐ False

Answer: MYALEPT is available only through pharmacies that are specially certified and agree to follow REMS requirements. For a list of certified pharmacies call: 1-855-669-2537.

Completion of Training for the MYALEPT™ (metreleptin) for injection REMS Program

You have completed training for the MYALEPT REMS Program.

To enroll in the MYALEPT REMS Program, complete the Enrollment Form and return via fax at 1-877-328-9682.

For more information on the MYALEPT REMS Program, please call 1-855-669-2537 or visit www.MYALEPTREMS.com.



©2014 AstraZeneca. All rights reserved.

MYALEPT is a trademark of the AstraZeneca group of companies.

MYALEPT is manufactured for Amylin Pharmaceuticals and distributed by AstraZeneca Pharmaceuticals LP.



i 1

MYALEPT REMS Program: An Introduction

What is the MYALEPT REMS (Risk Evaluation and Mitigation Strategy) Program?

A REMS is a strategy to manage known or potential risks associated with a drug, and is required by the FDA to ensure that the benefits of the drug outweigh its risks. MYALEPT is available only under a restricted program called the MYALEPT REMS Program because of:

- the development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT and the serious adverse events that may result
- · the risk of lymphoma.

Because of these risks, appropriate patient selection consistent with the approved indication for MYALEPT is very important.

MYALEPT REMS Program Requirements

- Certification of Prescribers of MYALEPT
- Completion of a Prescription Authorization Form for each new prescription
- Restricted distribution of MYALEPT through certified pharmacies

Certification of Prescribers of MYALEPT

- 1. Review the Prescribing Information and MYALEPT Prescriber Training Module
- 2. Complete, sign, and submit the one-time MYALEPT REMS Program Prescriber Enrollment Form

All Materials can be downloaded from the MYALEPT REMS website at: www.MYALEPTREMS.com. Or request these materials by calling 1-855-669-2537.

Completion of Prescription Authorization Form

Each new prescription for MYALEPT must be written using the MYALEPT Prescription Authorization Form.

- Download the Prescription Authorization Form at www.MYALEPTREMS.com or request a copy by calling 1-855-669-2537
- Complete the Prescription Authorization Form

Restricted Distribution of MYALEPT through Certified Pharmacies

Prescription Authorization Form must be signed and submitted by:

Fax to 1-877-328-9682





11.3 mg per vial



MYALEPT™ REMS Program

Program Requirements | Training & Enrollment

A Risk Evaluation and Mitigation Strategy (REMS) is a strategy to manage known or potential serious risks associated with a drug product and is required by the Food and Drug Administration (FDA) to ensure that the benefits of a drug outweigh its risks.

AstraZeneca has worked with the FDA to develop the MYALEPT REMS Program:

To educate prescribers about

- the development of anti-metreleptin antibodies that neutralize endogenous leptin and/or MYALEPT and the serious adverse events that may result from these antibodies.
- the risk of lymphoma, and
- appropriate patient selection

Program Requirements

MYALEPT is available only through the MYALEPT REMS Program. The MYALEPT REMS Program requirements include:

For Prescribers:

- Certification of prescribers of MYALEPT
 - Certification consists of completion of training, and enrollment in the MYALEPT REMS Program
- Completion of a Prescription Authorization form for each new prescription

Find out more about Training & Enrollment:

For Pharmacies:

 Restricted distribution of MYALEPT to patients with completed Prescription Authorization Forms from prescribers who are certified in the MYALEPT REMS Program

Training & Enrollment

Healthcare providers who prescribe MYALEPT must review the prescriber training materials to enroll in the MYALEPT REMS Program.

Steps to Prescriber Certification

- 1. Review the Prescriber Education Materials
 - MYALEPT Prescribing Information
 - Prescriber Training Module
- 2. Complete and submit the MYALEPT REMS Program Prescriber Enrollment Form
 - Print and sign the Prescriber Enrollment Form or request a copy by calling 1-855-669-2537
 - Submit the form via:
 - Fax to 1-877-328-9682

By completing the Prescriber Enrollment Form, the prescriber agrees to comply with the MYALEPT REMS Program requirements. A confirmation of your certification in the MYALEPT REMS program will be sent to you so you can begin to prescribe MYALEPT.

Reporting Adverse Reactions

Healthcare providers should report all suspected adverse events.

Please contact the company at 1-855-669-2537 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.com.

REMS Materials

- MYALEPT REMS Program:
 An Introduction
- MYALEPT REMS Program
 Prescriber Training Module
- MYALEPT REMS Program
 Prescriber Enrollment Form
- MYALEPT REMS Program
 Prescription Authorization Form
- Prescribing Information

MYALEPT REMS Program Contact Us

Phone: 1-855-669-2537

Fax: 1-877-328-9682

Hours of Operation:

Monday - Friday,
8:00am-8:00pm

Eastern Time

US Full Prescribing Information | Medication Guide | Important Safety Information | Privacy Notice | Legal Statement



©2014 AstraZeneca. All Rights Reserved.

MYALEPT is a trademark of the AstraZeneca group of companies.

MYALEPT is manufactured for Amylin Pharmaceuticals and distributed by AstraZeneca Pharmaceuticals LP.

1775310-2995302 Last Updated 6/14